

Referentielijst Organisatie van de zorg voor depressie

AkwaGGZ. (2017). Generieke module 'Arbeid als medicijn'. In. Utrecht: Akwa GGZ.

Archer, J., Bower, P., Gilbody, S., Lovell, K., Richards, D., Gask, L., Dickens, C., & Coventry, P. (2012). Collaborative care for depression and anxiety problems. The Cochrane Database of Systematic Reviews, 10, CD006525. <https://doi.org/10.1002/14651858.CD006525.pub2>

Bakker P, Jansen P. (2013). Generalistische Basis GGZ: verwijzmodel en productbeschrijvingen. Enschede: Bureau HHM.

Barkham, M., Rees, A., Stiles, W. B., Shapiro, D. A., Hardy, G. E., & Reynolds, S. (1996). Dose-effect relations in time-limited psychotherapy for depression. Journal of Consulting and Clinical Psychology, 64(5), 927–935. <https://doi.org/10.1037//0022-006x.64.5.927>

Borg, M., Karlsson, B., Lofthus, A. M., & Davidson, L. (2011). "Hitting the wall": Lived experiences of mental health crises. International journal of qualitative studies on health and well-being, 6(4), 10.3402/qhw.v6i4.7197. <https://doi.org/10.3402/qhw.v6i4.7197>

Bransen, E., Paalvast, M., & Slaats, M. (2014). De kracht van naasten ontsluiten. Tijdschrift Voor Psychotherapie, 40(1), 20–33.

Burns, A., Banerjee, S., Morris, J., Woodward, Y., Baldwin, R., Proctor, R., Tarrier, N., Pendleton, N., Sutherland, D., Andrew, G., & Horan, M. (2007). Treatment and prevention of depression after surgery for hip fracture in older people: randomized, controlled trials. Journal of the American Geriatrics Society, 55(1), 75–80. <https://doi.org/10.1111/j.1532-5415.2007.01016.x>

Canacott, L., Moghaddam, N., & Tickle, A. (2019). Is the Wellness Recovery Action Plan (WRAP) efficacious for improving personal and clinical recovery outcomes? A systematic review and meta-analysis. Psychiatric Rehabilitation Journal, 42(4), 372–381. <https://doi.org/10.1037/prj0000368>

Canacott, L., Tickle, A., & Moghaddam, N. (2020). Perceptions of wellness recovery action plan (WRAP) training: a systematic review and metasynthesis. Mental Health Review Journal, 25(4), 345–366. <https://doi.org/10.1108/MHRJ-10-2019-0037>

Cape, J., Whittington, C., & Bower, P. (2010). What is the role of consultation-liaison psychiatry in the management of depression in primary care? A systematic review and meta-analysis. General Hospital Psychiatry, 32(3), 246–254. <https://doi.org/10.1016/j.genhosppsych.2010.02.003>

Delgado J, et al. 2021. Cost-effectiveness of feedback-informed psychological treatment: Evidence from the IAPT-FIT trial. Behaviour Research and Therapy. <https://doi.org/10.1016/j.brat.2021.103873>

Dinger, U., Klipsch, O., Köhling, J., Ehrental, J. C., Nikendei, C., Herzog, W., & Schauenburg, H. (2014). Day-clinic and inpatient psychotherapy for depression (DIP-D): a randomized controlled pilot study in routine clinical care. In Psychotherapy and psychosomatics (Vol. 83, Issue 3, pp. 194–195). <https://doi.org/10.1159/000357437>

Gezondheidsraad. Verzekeringsgeneeskundige protocollen: Algemene inleiding, Overspanning, Depressieve stoornis. Den Haag: Gezondheidsraad, 2006; publicatie nr 2006/22.

Gilbody, S., Bower, P., Fletcher, J., Richards, D., & Sutton, A. J. (2006). Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. *Archives of Internal Medicine*, 166(21), 2314–2321. <https://doi.org/10.1001/archinte.166.21.2314>

Huijbregts, K. M., de Jong, F. J., van Marwijk, H. W., Beekman, A. T., Adèr, H. J., Hakkaart-van Roijen, L., Unützer, J., & van der Feltz-Cornelis, C. M. (2013). A target-driven collaborative care model for Major Depressive Disorder is effective in primary care in the Netherlands. A randomized clinical trial from the depression initiative. *Journal of affective disorders*, 146(3), 328–337. <https://doi.org/10.1016/j.jad.2012.09.015>

KNMG. (2017). KNMG-visiedocument Zorg die werkt. In *Naar een betere arbeidsgerichte medische zorg voor (potentieel) werkenden*. Utrecht: KNMG.

Meeuwissen, J.A.C. (2018). The case for stepped care - Exploring the applicability and cost-utility of stepped-care strategies in the management of depression. Amsterdam: VU University.

Meeuwissen, J.A.C., Feenstra, T.L., Smit, F., Blankers, M., Spijker, J., Bockting, C.L.H., van Balkom, A.J.L.M. & Buskens, E. (2019). The cost-utility of stepped-care algorithms according to depression guideline recommendations – results of a state-transition model analysis. *Journal of Affective Disorders*, Vol. 242, 244-254. <https://doi.org/10.1016/j.jad.2018.08.024>.

Mueller, M., & Pekarik, G. (2000). Treatment duration prediction: Client accuracy and its relationship to dropout, outcome, and satisfaction. *Psychotherapy: Theory, Research, Practice, Training*, 37, 117–123. <https://doi.org/10.1037/h0087701>

NVvP. (2017). Geestelijke gezondheid voor alle ouderen. Visiedocument november 2017. Geraadpleegd op 13 november 2023, van <https://www.trimbos.nl/docs/a1af681f-5b9b-496e-a72d-ceba87c43193.pdf>

Oosterbaan, D. B., Verbraak, M. J. P. M., Terluin, B., Hoogendoorn, A. W., Peyrot, W. J., Muntingh, A., & van Balkom, A. J. L. M. (2013). Collaborative stepped care v. care as usual for common mental disorders: 8-month, cluster randomised controlled trial. *The British Journal of Psychiatry : The Journal of Mental Science*, 203(2), 132–139. <https://doi.org/10.1192/bjp.bp.112.125211>

Owen, J., Smith, A., & Rodolfa, E. (2009). Clients' Expected Number of Counseling Sessions, Treatment Effectiveness, and Termination Status: Using Empirical Evidence to Inform Session Limit Policies. *Journal of College Student Psychotherapy*, 23(2), 118–134. <https://doi.org/10.1080/87568220902743660>

Peeters, F. P. M. L., Ruhe, H. G., Wichers, M., Abidi, L., Kaub, K., van der Lande, H. J., Spijker, J., Huibers, M. J. H., & Schene, A. H. (2016). The Dutch Measure for quantification of Treatment Resistance in Depression (DM-TRD): an extension of the Maudsley Staging Method. *Journal of Affective Disorders*, 205, 365–371. <https://doi.org/10.1016/j.jad.2016.08.019>

Penterman, E.J.M., Nijman, H.L.I. (2009). Het inschatten van agressie bij patiënten van de GGZ-crisisdienst. *Tijdschrift voor Psychiatrie*, 51, 6, (2009), pp. 355-364.

Repper, J., & Carter, T. (2011). A review of the literature on peer support in mental health services. *Journal of Mental Health (Abingdon, England)*, 20(4), 392–411. <https://doi.org/10.3109/09638237.2011.583947>

Reupert, A., Maybery, D., Nicholson, J., Gopfert, M., & Seeman, M. V (Eds.). (2015). Parental Psychiatric Disorder. In *Distressed Parents and their Families*. Cambridge University Press. <https://doi.org/10.1017/CBO9781107707559>

Rivero-Santana A, et al. (2021). Stepped care for the treatment of depression: a systematic review and meta-analysis. *Journal of Affective Disorders*. <https://doi.org/10.1016/j.jad.2021.07.008>

Schene, A., B Sabbe, Philip Spinhoven, Eric Ruhé (red.). (2016). *Handboek depressieve stoornissen*. Utrecht: De Tijdstroom.

Spijker, J., Bockting, C.L.H., Meeuwissen, J.A.C., Vliet, I.M. van, Emmelkamp, P.M.G., Hermens, M.L.M. & van Balkom A.L.J.M., namens de Werkgroep Multidisciplinaire richtlijnontwikkeling Angststoornissen/Depressie (2013). *Multidisciplinaire richtlijn Depressie (Derde revisie). Richtlijn voor de diagnostiek, behandeling en begeleiding van volwassen patiënten met een depressieve stoornis*. Utrecht: Trimbos-instituut.

Sytema, S., Wunderink, L., Bloemers, W., Roorda, L., & Wiersma, D. (2007). Assertive community treatment in the Netherlands: a randomized controlled trial. *Acta psychiatrica Scandinavica*, 116(2), 105–112. <https://doi.org/10.1111/j.1600-0447.2007.01021.x>

Tjaden, C., Mulder, C. L., den Hollander, W., Castelein, S., Delespaul, P., Keet, R., van Weeghel, J., & Kroon, H. (2021). Effectiveness of Resource Groups for Improving Empowerment, Quality of Life, and Functioning of People With Severe Mental Illness: A Randomized Clinical Trial. *JAMA Psychiatry*, 78(12), 1309–1318. <https://doi.org/10.1001/jamapsychiatry.2021.2880>

UWV. (2023). *Ik ben ziek (WIA-uitkering)*. <https://www.uwv.nl/particulieren/ziek/ziek-wia-uitkering/index.aspx>

Van der Feltz-Cornelis, C. M., Nuyen, J., Stoop, C., Chan, J., Jacobson, A. M., Katon, W., Snoek, F., & Sartorius, N. (2010). Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a systematic review and meta-analysis. *General hospital psychiatry*, 32(4), 380–395. <https://doi.org/10.1016/j.genhosppsych.2010.03.011>

Van Alphen, C., Ammeraal, M., Blanke, C., Boonstra, N., Boumans, H., Bruggeman, R., Castelein, S., Dekker, F.L., van Duin, D., van Ewijk, W.M., van der Gaag, M., van Gool, R., de Haas, O., Henquet, C., Hermens, M.L.M., Ketelaars, T., Knegtering, H., Krans, M.J., Lansens, M., Lochy, R., Loonen, A.J.M., Meesters, P.D., van Meijel, B.K.G., Miltenburg, G., Oud, M.J.T., de Pater, C.F., Peterse, A., van der Plas, J., Rammers, D., Rentenaar, I.M.F., van Santen, G.W., Scheewe, T.W., Selten, J.P., Slooff, C.J.A.J., Stoop, C., Vlamincck, P., Vollaard, H., van Weeghel, J., van Wel, T.. (2012). *Multidisciplinaire richtlijn schizofrenie*. Utrecht: De Tijdstroom.

Van der Veen, C., Voordouw, I., & Harmsen, K. (2008). *Aard en omvang GGZ- en VZ-preventie 2006; ontwikkelingen in de organisatie, omvang en inhoud van de GGZ- en VZ-preventie*.

Van der Voort, T. Y. G., Goossens, P. J. J., & van der Bijl, J. J. (2009). Alone together: A grounded theory study of experienced burdens, coping, and support needs of spouses of persons with a bipolar disorder. *International Journal of Mental Health Nursing*, 18(6), 434–443. <https://doi.org/10.1111/j.1447-0349.2009.00634.x>

- Van Dijk, D. A., van den Boogaard, T. M., Deen, M. L., Spijker, J., Ruhé, H. G., & Peeters, F. P. M. L. (2019). Predicting clinical course in major depressive disorder: The association between DM-TRD score and symptom severity over time in 1115 outpatients. *Depression and Anxiety*, 36(4), 345–352. <https://doi.org/10.1002/da.22865>
- Van Gestel-Timmermans, H., Brouwers, E. P. M., van Assen, M. A. L. M., & van Nieuwenhuizen, C. (2012). Effects of a peer-run course on recovery from serious mental illness: a randomized controlled trial. *Psychiatric Services (Washington, D.C.)*, 63(1), 54–60. <https://doi.org/10.1176/appi.ps.201000450>
- Van Krugten, F. C. W., Goorden, M., van Balkom, A. J. L. M., van Oppen, P., Ruhé, H. G., van Schaik, D. J. F., Brouwer, W. B. F., & Hakkaart-van Roijen, L. (2019). The decision tool unipolar depression (DTUD): a new measure to facilitate the early identification of patients with major depressive disorder in need of highly specialized care. *BMC Psychiatry*, 19(1), 179. <https://doi.org/10.1186/s12888-019-2165-9>
- Van Wijngaarden, B. (2003). Consequences for caregivers of patients with severe Mental illness: the development of the involvement evaluation questionnaire. Universiteit van Amsterdam.
- Visser E, Kregting B, Olde Hartman T, Pop-Purceanu M, van de Laar F. (2023). Positieve ervaringen consultatieve psychiatrie in spreekkamer. *Huisarts Wet*; 66:DOI:10.1007/s12445-023-2274-z.
- White, S., Foster, R., Marks, J., Morshead, R., Goldsmith, L., Barlow, S., Sin, J., & Gillard, S. (2020). The effectiveness of one-to-one peer support in mental health services: a systematic review and meta-analysis. *BMC Psychiatry*, 20(1), 534. <https://doi.org/10.1186/s12888-020-02923-3>
- Zeeck, A., von Wietersheim, J., Weiß, H., Eduard Scheidt, C., Völker, A., Helesic, A., Eckhardt-Henn, A., Beutel, M., Endorf, K., Knoblauch, J., Rochlitz, P., & Hartmann, A. (2015). Symptom course in inpatient and day clinic treatment of depression: Results from the INDDEP-Study. *Journal of Affective Disorders*, 187, 35–44. <https://doi.org/10.1016/j.jad.2015.07.025>
- Zhu, M., Ran Ha Hong, Tao Yang, Xiaorui Yang, Xing Wang, Jing Liu, Jill K Murphy, Erin E Michalak, Zuowei Wang, Lakshmi N Yatham, Jun Chen, Raymond W Lam. (2021). The Efficacy of Measurement-Based Care for Depressive Disorders: Systematic Review and Meta-Analysis of Randomized Controlled Trials. *J Clin Psychiatry*. Sep 28;82(5):21r14034. doi: 10.4088/JCP.21r14034.