

Tabel 1 Overzicht van enkele studies die contact allergenen hebben onderzocht

| Author | Origin leg ulcer | Study type | N patients | N allergens tested | N (%) patients with sensitization to at least 1 allergen | Top 3 allergens, N (%) | Top 3 wound dressings allergens, N (%) | Correlation and/or remarks |
|-----------------------------------|--|--------------------------|------------|--------------------|--|---|--|---|
| Malten KE, Kuiper JP (1985) | VLU | Retrospective | 100 | 92 | 69 (69) ≥2, 57 (57) | Lanolin, 17 (17) Fragrance, 14 (14) PPD, 9 (9) Wood tar, 9 (9) | | |
| Tavadia et al (2003) | VLU | Observation | 200 | 90 | 136 (68) ≥2, 102 (51) ≥5, 36 (18) | Fragrance, 47 (23,5) Lanolin alcohol, 33 (16,5) Balsam of Peru, 29 (14,5) Aminoglycosides, 29 (14,5) | Intrasite gel, 19 (9,5) Hioxyl cream, 17 (8,5) Trimovate cream, 13 (6,5) | Remark: control group was tested, no % are stated |
| Machet et al (2006) | leg ulcer + Meta-analysis data 1973-2003 | Retrospective | 106 | 36 | 80 (75) ≥2, 60 (57) | Balsam of Peru, 42 (39,6) Lanolin, 22 (20,8) Fragrance, 19 (17,9) | Triclocarban, 14 (13,2) Cetavlon cream, 10 (9,4) Betadine cream, 8 (7,5) | Remark: control group eczema sensitization 54% |
| Smart et al (2008) | VLU and AV leg ulcer | Prospective | 100 | 38 | 46 (46) ≥2, 39 (39) | Wood tar, 21 (21) Lanolin alcohol, 11 (11) Balsam of Peru, 10 (10) | | |
| Barbaud et al (2009) | Chronic: 55% VLU | Prospective multi-centre | 423 | 62 | 308 (73) Antiseptics, 85 (20) | Balsam of Peru, 172 (40) Fragrance, 112 (26,5) Antiseptics, 85 (20,1) | Biafine, 36 (8,5) Duoderm E, 17 (4) Comfeel transparent, 6 (1,4) | + duration ulcer, + eczema (even if 57% +) |

| | | | | | | | | |
|-----------------------|----------------------------------|-----------------------------|-----|----|------------------------------------|--|--|--|
| | | | | | | | | reaction without eczema) No relation with cause |
| Valois et al (2015) | Chronic: 54% VLU 20% AV | Prospective multi-centre | 354 | 64 | 211 (59,6) ≥ 2 , 90 (25,4) | Balsam of Peru, 84 (23,7) Fragrance, 47 (13,3) Benzalkonium- chloride, 25 (7) | Ialuset cream, 45 (12,7) Hydrocellular, 28 (7,9) Duoderm E, 18 (5,1) | + duration leg ulcer history No relation with cause, eczema or duration present ulcer |
| Rai et al (2018) | VLU India | Prospective multi-centre | 172 | 27 | 83 (48,2) ≥ 2 , 59 (34,3) | Wood tar, 19 (10,4) Framycetin, 16 (8,7) Thiomersal, 13 (7,1) Eosin, 13 (7,1) | | Not studied |
| Raudonis et al (2019) | Chronic: 65% VLU | Retrospective | 145 | 68 | 91 (62,8) | Benzalkonium chl, 27 (18,6) Balsam of Peru, 26 (17,9) Povidone-iodine, 25 (17,2) | | + with pruritus/eczema |