

Tabel 2. Uitgangsvraag 2 - Welke klachten komen voor tijdens de (ziekte- en symptoomgerichte) palliatieve fase?

Author year	Design	Selection criteria	Quality	Results	Level of Evidence
Solano 2006	Systematic review	Studies reporting symptom prevalence in patients with advanced illness suffering from AIDS, heart disease, COPD, renal disease or cancer	Search +, Selection +, Quality ass ±	64 articles included. Depression common in all chronic diseases (60-82% maximum prevalences). Constipation and anorexia most common in cancer patients	A2
Teunissen 2007b	Systematic review	Studies reporting symptom prevalence in patients with incurable cancer	Search ±, Selection +, Quality ass ?, Pooling +	46 studies on 26223 patients included. Pain, lack of energy, weakness and appetite loss occurred in >50%. Among patients in last 1-2 weeks of life occurred fatigue, weight loss, weakness and appetite loss in >50%	A2
Van den Beuken 2007	Systematic review	Studies reporting prevalence of pain in patients with cancer	Search +, Selection +, Quality ass +, Pooling +	52 studies included. Prevalence of pain 64% (CI 58-69%) in patients with advanced disease; 59% (CI 44-73%) in patients on anti-cancer treatment; 33% (CI 21-46%) after curative treatment.	A2